

Bounds on the cardinality of partition

Kerry M. Soileau

November 14, 2007

ABSTRACT

If A is infinite and well-ordered, then $|2^A| \leq |Part(A)| \leq |A^A|$.

INTRODUCTION

A partition of a set A is a collection of nonempty disjoint subsets of A whose union is

A . Let $Part(A)$ be the collection of partitions of a set A .

Theorem: If A is infinite and well-ordered, then $|2^A| \leq |Part(A)| \leq |A^A|$

For each nonempty subset $P \subseteq A$ define $s(P) = \min(m \in P)$ under the well-ordering on A .

Proof of $|2^A| \leq |Part(A)|$: Define $f : 2^A \rightarrow Part(A)$ verifying

$$f(S) = \begin{cases} \{A\} & \text{if } S = \emptyset \\ \{\{s(A)\}, A \sim \{s(A)\}\} & \text{if } S = A \\ \{\{s(S)\}, S \sim \{s(S)\}, A \sim S\} & \text{otherwise} \end{cases}$$

Now suppose $f(S_1) = f(S_2)$. If $|f(S_1)| = 1$, then $S_1 = S_2 = \emptyset$ so $S_1 = S_2$. If $|f(S_1)| = 2$, then $S_1 = S_2 = A$ so $S_1 = S_2$. If $|f(S_1)| = 3$, then clearly $A \sim S_1 = A \sim S_2$, hence $S_1 = S_2$ and thus f is an injection.

Proof of $|Part(A)| \leq |A^A|$: Define $g : Part(A) \rightarrow A^A$ verifying

$\{g(\pi)(a)\} = \{s(P); P \in \pi \text{ and } a \in P\}$ for $\pi \in Part(A)$ and $a \in A$. Then g is an injection.

REFERENCES

1. Cantor, Georg (1955, 1915). Contributions to the Founding of the Theory of Transfinite Numbers. New York: Dover.